

表 734 心筋梗塞の既往 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	51,873	4,158	56,031	178	12,368	68,577
(%)	(92.6)	(7.4)	(100.0)			
慢性腎盂腎炎	1,700	107	1,807	5	353	2,165
(%)	(94.1)	(5.9)	(100.0)			
急速進行性糸球体腎炎	1,433	95	1,528	7	299	1,834
(%)	(93.8)	(6.2)	(100.0)			
妊娠腎/妊娠中毒症	848	42	890	5	169	1,064
(%)	(95.3)	(4.7)	(100.0)			
その他の分類不能の腎炎	722	39	761		188	949
(%)	(94.9)	(5.1)	(100.0)			
多発性嚢胞腎	6,354	410	6,764	20	1,414	8,198
(%)	(93.9)	(6.1)	(100.0)			
腎硬化症	16,189	1,861	18,050	96	3,314	21,460
(%)	(89.7)	(10.3)	(100.0)			
悪性高血圧	1,435	107	1,542	3	336	1,881
(%)	(93.1)	(6.9)	(100.0)			
糖尿病性腎症	64,358	10,102	74,460	307	15,740	90,507
(%)	(86.4)	(13.6)	(100.0)			
SLE腎炎	1,226	76	1,302	4	244	1,550
(%)	(94.2)	(5.8)	(100.0)			
アミロイド腎	248	10	258	3	64	325
(%)	(96.1)	(3.9)	(100.0)			
痛風腎	654	48	702	2	124	828
(%)	(93.2)	(6.8)	(100.0)			
先天性代謝異常による腎不全	142	7	149		28	177
(%)	(95.3)	(4.7)	(100.0)			
腎・尿路結核	119	10	129		29	158
(%)	(92.2)	(7.8)	(100.0)			
腎・尿路結石	328	23	351		79	430
(%)	(93.4)	(6.6)	(100.0)			
腎・尿路腫瘍	518	32	550	2	134	686
(%)	(94.2)	(5.8)	(100.0)			
閉塞性尿路障害	417	23	440	3	74	517
(%)	(94.8)	(5.2)	(100.0)			
骨髄腫	152	13	165	1	24	190
(%)	(92.1)	(7.9)	(100.0)			
腎形成不全	343	10	353		66	419
(%)	(97.2)	(2.8)	(100.0)			
不明	14,833	1,321	16,154	60	3,911	20,125
(%)	(91.8)	(8.2)	(100.0)			
再導入	1,090	79	1,169		248	1,417
(%)	(93.2)	(6.8)	(100.0)			
その他	4,165	359	4,524	30	960	5,514
(%)	(92.1)	(7.9)	(100.0)			
合計	169,147	18,932	188,079	726	40,166	228,971
(%)	(89.9)	(10.1)	(100.0)			
記載なし						
(%)						
総計	169,147	18,932	188,079	726	40,166	228,971
(%)	(89.9)	(10.1)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査