

表 745 心筋梗塞の既往 血流量(ml/分)別（施設血液透析患者 透析回数週3回）

血流量	既往なし	既往あり	合計	不明	記載なし	総計
100未満	34	4	38		6	44
(%)	(89.5)	(10.5)	(100.0)			
100～	580	72	652	3	84	739
(%)	(89.0)	(11.0)	(100.0)			
120～	1,644	206	1,850	18	280	2,148
(%)	(88.9)	(11.1)	(100.0)			
140～	13,502	1,620	15,122	65	2,786	17,973
(%)	(89.3)	(10.7)	(100.0)			
160～	4,596	545	5,141	12	748	5,901
(%)	(89.4)	(10.6)	(100.0)			
180～	22,915	2,635	25,550	134	4,801	30,485
(%)	(89.7)	(10.3)	(100.0)			
200～	74,978	8,749	83,727	226	16,537	100,490
(%)	(89.6)	(10.4)	(100.0)			
220～	21,848	2,323	24,171	91	5,284	29,546
(%)	(90.4)	(9.6)	(100.0)			
240～	20,411	1,941	22,352	73	4,635	27,060
(%)	(91.3)	(8.7)	(100.0)			
260～	1,884	185	2,069	8	456	2,533
(%)	(91.1)	(8.9)	(100.0)			
280～	1,751	163	1,914	2	421	2,337
(%)	(91.5)	(8.5)	(100.0)			
300～	3,178	248	3,426	5	834	4,265
(%)	(92.8)	(7.2)	(100.0)			
350～	352	18	370	1	58	429
(%)	(95.1)	(4.9)	(100.0)			
400～	467	38	505	1	212	718
(%)	(92.5)	(7.5)	(100.0)			
合計	168,140	18,747	186,887	639	37,142	224,668
(%)	(90.0)	(10.0)	(100.0)			
記載なし	1,007	185	1,192	87	3,024	4,303
(%)	(84.5)	(15.5)	(100.0)			
総計	169,147	18,932	188,079	726	40,166	228,971
(%)	(89.9)	(10.1)	(100.0)			
平均	204.74	202.00	204.46	198.85	207.42	204.94
標準偏差	36.07	33.97	35.87	35.64	38.95	36.42

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査