

表 746 心筋梗塞の既往 身長(cm)別（施設血液透析患者 透析回数週3回）

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	181	26	207	2	28	237
(%)	(87.4)	(12.6)	(100.0)			
130～	452	46	498		70	568
(%)	(90.8)	(9.2)	(100.0)			
135～	1,781	157	1,938	16	351	2,305
(%)	(91.9)	(8.1)	(100.0)			
140～	6,459	539	6,998	42	1,243	8,283
(%)	(92.3)	(7.7)	(100.0)			
145～	14,632	1,314	15,946	56	2,817	18,819
(%)	(91.8)	(8.2)	(100.0)			
150～	23,067	2,146	25,213	90	4,711	30,014
(%)	(91.5)	(8.5)	(100.0)			
155～	26,605	2,887	29,492	105	5,260	34,857
(%)	(90.2)	(9.8)	(100.0)			
160～	31,612	4,087	35,699	111	6,448	42,258
(%)	(88.6)	(11.4)	(100.0)			
165～	28,103	3,752	31,855	79	5,642	37,576
(%)	(88.2)	(11.8)	(100.0)			
170～	17,935	2,116	20,051	67	3,709	23,827
(%)	(89.4)	(10.6)	(100.0)			
175～	6,552	647	7,199	18	1,332	8,549
(%)	(91.0)	(9.0)	(100.0)			
180～	1,762	137	1,899	8	347	2,254
(%)	(92.8)	(7.2)	(100.0)			
185～	285	11	296		55	351
(%)	(96.3)	(3.7)	(100.0)			
190～	44	5	49		8	57
(%)	(89.8)	(10.2)	(100.0)			
合計	159,470	17,870	177,340	594	32,021	209,955
(%)	(89.9)	(10.1)	(100.0)			
記載なし	9,677	1,062	10,739	132	8,145	19,016
(%)	(90.1)	(9.9)	(100.0)			
総計	169,147	18,932	188,079	726	40,166	228,971
(%)	(89.9)	(10.1)	(100.0)			
平均	159.85	160.72	159.94	158.22	159.98	159.94
標準偏差	9.64	9.09	9.59	10.68	9.56	9.59

数値下のカッコ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査