

表 750 心筋梗塞の既往 Body Mass Index別（施設血液透析患者 透析回数週3回）

Body Mass Index	既往なし	既往あり	合計	不明	記載なし	総計
14未満	1,001	96	1,097	2	212	1,311
(%)	(91.2)	(8.8)	(100.0)			
14～	5,890	562	6,452	33	1,121	7,606
(%)	(91.3)	(8.7)	(100.0)			
16～	18,848	1,899	20,747	69	3,664	24,480
(%)	(90.8)	(9.2)	(100.0)			
18～	33,115	3,562	36,677	122	6,357	43,156
(%)	(90.3)	(9.7)	(100.0)			
20～	36,379	4,029	40,408	121	7,151	47,680
(%)	(90.0)	(10.0)	(100.0)			
22～	27,905	3,319	31,224	95	5,491	36,810
(%)	(89.4)	(10.6)	(100.0)			
24～	16,565	2,065	18,630	67	3,216	21,913
(%)	(88.9)	(11.1)	(100.0)			
26～	8,823	1,077	9,900	36	1,752	11,688
(%)	(89.1)	(10.9)	(100.0)			
28～	4,545	546	5,091	19	881	5,991
(%)	(89.3)	(10.7)	(100.0)			
30～	4,007	405	4,412	14	738	5,164
(%)	(90.8)	(9.2)	(100.0)			
合計	157,078	17,560	174,638	578	30,583	205,799
(%)	(89.9)	(10.1)	(100.0)			
記載なし	12,069	1,372	13,441	148	9,583	23,172
(%)	(89.8)	(10.2)	(100.0)			
総計	169,147	18,932	188,079	726	40,166	228,971
(%)	(89.9)	(10.1)	(100.0)			
平均	21.45	21.64	21.47	21.40	21.45	21.46
標準偏差	3.66	3.61	3.66	3.75	3.65	3.66

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査