

表 751 心筋梗塞の既往 透析前BUN(mg/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前BUN	既往なし	既往あり	合計	不明	記載なし	総計
20未満	355	38	393	4	82	479
(%)	(90.3)	(9.7)	(100.0)			
20～	1,806	253	2,059	9	412	2,480
(%)	(87.7)	(12.3)	(100.0)			
30～	7,484	1,113	8,597	27	1,734	10,358
(%)	(87.1)	(12.9)	(100.0)			
40～	19,646	2,737	22,383	82	4,346	26,811
(%)	(87.8)	(12.2)	(100.0)			
50～	31,523	3,992	35,515	103	7,041	42,659
(%)	(88.8)	(11.2)	(100.0)			
60～	31,848	3,643	35,491	99	7,053	42,643
(%)	(89.7)	(10.3)	(100.0)			
70～	21,391	2,202	23,593	78	4,651	28,322
(%)	(90.7)	(9.3)	(100.0)			
80～	10,048	949	10,997	35	2,180	13,212
(%)	(91.4)	(8.6)	(100.0)			
90～	3,611	359	3,970	12	821	4,803
(%)	(91.0)	(9.0)	(100.0)			
100～	1,057	83	1,140	8	200	1,348
(%)	(92.7)	(7.3)	(100.0)			
110～	304	26	330		64	394
(%)	(92.1)	(7.9)	(100.0)			
120～	150	18	168	1	35	204
(%)	(89.3)	(10.7)	(100.0)			
合計	129,223	15,413	144,636	458	28,619	173,713
(%)	(89.3)	(10.7)	(100.0)			
記載なし	1,426	235	1,661	54	2,781	4,496
(%)	(85.9)	(14.1)	(100.0)			
総計	130,649	15,648	146,297	512	31,400	178,209
(%)	(89.3)	(10.7)	(100.0)			
平均	61.15	59.16	60.94	60.48	60.95	60.94
標準偏差	15.65	15.41	15.64	16.83	15.63	15.64

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査