

表 762 心筋梗塞の既往 PTH値(pg/mL)別（施設血液透析患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	1,142	137	1,279	2	218	1,499
(%)	(89.3)	(10.7)	(100.0)			
20～	1,443	179	1,622	2	224	1,848
(%)	(89.0)	(11.0)	(100.0)			
35～	1,236	154	1,390		223	1,613
(%)	(88.9)	(11.1)	(100.0)			
50～	1,888	229	2,117	3	339	2,459
(%)	(89.2)	(10.8)	(100.0)			
75～	1,629	197	1,826	3	319	2,148
(%)	(89.2)	(10.8)	(100.0)			
100～	1,353	150	1,503	4	280	1,787
(%)	(90.0)	(10.0)	(100.0)			
125～	870	96	966	1	220	1,187
(%)	(90.1)	(9.9)	(100.0)			
150～	30	2	32		5	37
(%)	(93.8)	(6.3)	(100.0)			
151～	708	80	788		146	934
(%)	(89.8)	(10.2)	(100.0)			
180～	502	60	562		104	666
(%)	(89.3)	(10.7)	(100.0)			
210～	344	40	384		88	472
(%)	(89.6)	(10.4)	(100.0)			
240～	402	37	439		87	526
(%)	(91.6)	(8.4)	(100.0)			
300～	159	7	166		39	205
(%)	(95.8)	(4.2)	(100.0)			
350～	100	14	114		21	135
(%)	(87.7)	(12.3)	(100.0)			
400～	123	8	131		21	152
(%)	(93.9)	(6.1)	(100.0)			
500～	128	13	141		25	166
(%)	(90.8)	(9.2)	(100.0)			
合計	12,057	1,403	13,460	15	2,359	15,834
(%)	(89.6)	(10.4)	(100.0)			
記載なし	233	23	256		182	438
(%)	(91.0)	(9.0)	(100.0)			
総計	12,290	1,426	13,716	15	2,541	16,272
(%)	(89.6)	(10.4)	(100.0)			
平均	105.40	98.56	104.69	72.73	111.66	105.70
標準偏差	110.89	99.45	109.77	40.57	106.29	109.24

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査