

表 763 心筋梗塞の既往 透析前ヘモグロビン濃度(g/dL)別（施設血液透析患者 透析回数週3回）

透析前ヘモグロビン濃度	既往なし	既往あり	合計	不明	記載なし	総計
5未満	208	24	232		29	261
(%)	(89.7)	(10.3)	(100.0)			
5～	285	25	310	1	51	362
(%)	(91.9)	(8.1)	(100.0)			
6～	687	74	761	15	166	942
(%)	(90.3)	(9.7)	(100.0)			
7～	2,687	304	2,991	18	569	3,578
(%)	(89.8)	(10.2)	(100.0)			
8～	8,855	1,055	9,910	43	1,773	11,726
(%)	(89.4)	(10.6)	(100.0)			
9～	27,866	3,193	31,059	103	5,876	37,038
(%)	(89.7)	(10.3)	(100.0)			
10～	56,768	6,160	62,928	222	12,067	75,217
(%)	(90.2)	(9.8)	(100.0)			
11～	46,395	5,081	51,476	145	9,858	61,479
(%)	(90.1)	(9.9)	(100.0)			
12～	16,811	1,910	18,721	62	3,587	22,370
(%)	(89.8)	(10.2)	(100.0)			
13～	4,130	509	4,639	7	891	5,537
(%)	(89.0)	(11.0)	(100.0)			
14～	1,445	210	1,655	4	274	1,933
(%)	(87.3)	(12.7)	(100.0)			
合計	166,137	18,545	184,682	620	35,141	220,443
(%)	(90.0)	(10.0)	(100.0)			
記載なし	3,010	387	3,397	106	5,025	8,528
(%)	(88.6)	(11.4)	(100.0)			
総計	169,147	18,932	188,079	726	40,166	228,971
(%)	(89.9)	(10.1)	(100.0)			
平均	10.69	10.70	10.69	10.46	10.70	10.69
標準偏差	1.28	1.31	1.28	1.39	1.26	1.28

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査