

表 769 心筋梗塞の既往 透析前脈拍(拍/分)別 (施設血液透析患者 透析回数週3回)

透析前脈拍	既往なし	既往あり	合計	不明	記載なし	総計
40未満	110	21	131		24	155
(%)	(84.0)	(16.0)	(100.0)			
40～	1,423	217	1,640	2	287	1,929
(%)	(86.8)	(13.2)	(100.0)			
50～	11,939	1,479	13,418	51	2,304	15,773
(%)	(89.0)	(11.0)	(100.0)			
60～	36,968	4,298	41,266	129	7,346	48,741
(%)	(89.6)	(10.4)	(100.0)			
70～	44,237	4,888	49,125	159	8,900	58,184
(%)	(90.0)	(10.0)	(100.0)			
80～	27,694	2,976	30,670	117	5,936	36,723
(%)	(90.3)	(9.7)	(100.0)			
90～	12,074	1,289	13,363	50	2,505	15,918
(%)	(90.4)	(9.6)	(100.0)			
100～	3,752	420	4,172	15	717	4,904
(%)	(89.9)	(10.1)	(100.0)			
110～	894	85	979	4	174	1,157
(%)	(91.3)	(8.7)	(100.0)			
120～	392	43	435	2	61	498
(%)	(90.1)	(9.9)	(100.0)			
合計	139,483	15,716	155,199	529	28,254	183,982
(%)	(89.9)	(10.1)	(100.0)			
記載なし	29,664	3,216	32,880	197	11,912	44,989
(%)	(90.2)	(9.8)	(100.0)			
総計	169,147	18,932	188,079	726	40,166	228,971
(%)	(89.9)	(10.1)	(100.0)			
平均	74.52	73.88	74.45	75.22	74.73	74.50
標準偏差	12.82	12.85	12.83	13.00	12.63	12.80

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査