

表 785 心筋梗塞の既往 身長(cm)別（血液透析濾過患者 透析回数週3回）

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	30	3	33		5	38
(%)	(90.9)	(9.1)	(100.0)			
130～	46	2	48	2	20	70
(%)	(95.8)	(4.2)	(100.0)			
135～	217	23	240	3	46	289
(%)	(90.4)	(9.6)	(100.0)			
140～	889	84	973	3	187	1,163
(%)	(91.4)	(8.6)	(100.0)			
145～	2,234	238	2,472	4	519	2,995
(%)	(90.4)	(9.6)	(100.0)			
150～	4,006	427	4,433	8	771	5,212
(%)	(90.4)	(9.6)	(100.0)			
155～	4,641	519	5,160	12	907	6,079
(%)	(89.9)	(10.1)	(100.0)			
160～	5,332	762	6,094	6	1,077	7,177
(%)	(87.5)	(12.5)	(100.0)			
165～	5,042	684	5,726	7	989	6,722
(%)	(88.1)	(11.9)	(100.0)			
170～	3,625	444	4,069	6	710	4,785
(%)	(89.1)	(10.9)	(100.0)			
175～	1,381	135	1,516	1	269	1,786
(%)	(91.1)	(8.9)	(100.0)			
180～	391	34	425		68	493
(%)	(92.0)	(8.0)	(100.0)			
185～	58	8	66		11	77
(%)	(87.9)	(12.1)	(100.0)			
190～	8		8		1	9
(%)	(100.0)		(100.0)			
合計	27,900	3,363	31,263	52	5,580	36,895
(%)	(89.2)	(10.8)	(100.0)			
記載なし	1,260	127	1,387	20	1,644	3,051
(%)	(90.8)	(9.2)	(100.0)			
総計	29,160	3,490	32,650	72	7,224	39,946
(%)	(89.3)	(10.7)	(100.0)			
平均	160.75	161.25	160.80	156.31	160.42	160.74
標準偏差	9.54	9.13	9.50	11.29	9.72	9.54

数値下のカッコ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査