

表 786 心筋梗塞の既往 透析後体重(kg)別（血液透析濾過患者 透析回数週3回）

透析後体重	既往なし	既往あり	合計	不明	記載なし	総計
20未満	8		8		4	12
(%)	(100.0)		(100.0)			
20～	59	4	63		13	76
(%)	(93.7)	(6.3)	(100.0)			
30～	2,000	195	2,195	11	476	2,682
(%)	(91.1)	(8.9)	(100.0)			
40～	7,457	857	8,314	16	1,675	10,005
(%)	(89.7)	(10.3)	(100.0)			
50～	9,188	1,181	10,369	20	2,101	12,490
(%)	(88.6)	(11.4)	(100.0)			
60～	6,228	756	6,984	12	1,369	8,365
(%)	(89.2)	(10.8)	(100.0)			
70～	2,593	297	2,890	1	532	3,423
(%)	(89.7)	(10.3)	(100.0)			
80～	962	112	1,074	2	197	1,273
(%)	(89.6)	(10.4)	(100.0)			
90～	337	39	376		79	455
(%)	(89.6)	(10.4)	(100.0)			
100～	105	8	113		29	142
(%)	(92.9)	(7.1)	(100.0)			
110～	50	3	53		8	61
(%)	(94.3)	(5.7)	(100.0)			
120～	29	3	32		5	37
(%)	(90.6)	(9.4)	(100.0)			
合計	29,016	3,455	32,471	62	6,488	39,021
(%)	(89.4)	(10.6)	(100.0)			
記載なし	144	35	179	10	736	925
(%)	(80.4)	(19.6)	(100.0)			
総計	29,160	3,490	32,650	72	7,224	39,946
(%)	(89.3)	(10.7)	(100.0)			
平均	56.69	56.87	56.71	52.06	56.27	56.63
標準偏差	13.11	12.40	13.04	11.17	13.00	13.03

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査