

表 801 心筋梗塞の既往 PTH値(pg/mL)別（血液透析濾過患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	197	19	216		33	249
(%)	(91.2)	(8.8)	(100.0)			
20～	237	27	264		37	301
(%)	(89.8)	(10.2)	(100.0)			
35～	209	28	237		37	274
(%)	(88.2)	(11.8)	(100.0)			
50～	331	42	373		71	444
(%)	(88.7)	(11.3)	(100.0)			
75～	301	21	322	1	68	391
(%)	(93.5)	(6.5)	(100.0)			
100～	253	29	282		49	331
(%)	(89.7)	(10.3)	(100.0)			
125～	178	16	194		40	234
(%)	(91.8)	(8.2)	(100.0)			
150～	4		4			4
(%)	(100.0)		(100.0)			
151～	157	29	186		29	215
(%)	(84.4)	(15.6)	(100.0)			
180～	108	10	118		19	137
(%)	(91.5)	(8.5)	(100.0)			
210～	79	6	85		13	98
(%)	(92.9)	(7.1)	(100.0)			
240～	69	10	79		9	88
(%)	(87.3)	(12.7)	(100.0)			
300～	34	3	37		5	42
(%)	(91.9)	(8.1)	(100.0)			
350～	24	3	27		2	29
(%)	(88.9)	(11.1)	(100.0)			
400～	15	5	20		4	24
(%)	(75.0)	(25.0)	(100.0)			
500～	36	6	42		3	45
(%)	(85.7)	(14.3)	(100.0)			
合計	2,232	254	2,486	1	419	2,906
(%)	(89.8)	(10.2)	(100.0)			
記載なし	8	1	9		22	31
(%)	(88.9)	(11.1)	(100.0)			
総計	2,240	255	2,495	1	441	2,937
(%)	(89.8)	(10.2)	(100.0)			
平均	112.01	120.43	112.87	77.00	105.48	111.79
標準偏差	109.93	117.77	110.76		90.82	108.12

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査