

表 829 心筋梗塞の既往 リン濃度(mg/dL)別（腹膜透析患者）

リン濃度	既往なし	既往あり	合計	不明	記載なし	総計
3未満	97	11	108	1	17	126
(%)	(89.8)	(10.2)	(100.0)			
3.0～	210	28	238	2	30	270
(%)	(88.2)	(11.8)	(100.0)			
3.5～	417	32	449	6	70	525
(%)	(92.9)	(7.1)	(100.0)			
4.0～	606	56	662	6	102	770
(%)	(91.5)	(8.5)	(100.0)			
4.5～	746	73	819	4	98	921
(%)	(91.1)	(8.9)	(100.0)			
5.0～	751	66	817	4	127	948
(%)	(91.9)	(8.1)	(100.0)			
5.5～	678	52	730	3	106	839
(%)	(92.9)	(7.1)	(100.0)			
6.0～	114	6	120		18	138
(%)	(95.0)	(5.0)	(100.0)			
6.1～	362	42	404	5	68	477
(%)	(89.6)	(10.4)	(100.0)			
6.5～	299	28	327	6	66	399
(%)	(91.4)	(8.6)	(100.0)			
7.0～	170	11	181	2	29	212
(%)	(93.9)	(6.1)	(100.0)			
7.5～	116	9	125	5	19	149
(%)	(92.8)	(7.2)	(100.0)			
8.0～	50	6	56	2	13	71
(%)	(89.3)	(10.7)	(100.0)			
8.5～	29	2	31		7	38
(%)	(93.5)	(6.5)	(100.0)			
9.0～	66	5	71	1	17	89
(%)	(93.0)	(7.0)	(100.0)			
合計	4,711	427	5,138	47	787	5,972
(%)	(91.7)	(8.3)	(100.0)			
記載なし	649	90	739	19	2,211	2,969
(%)	(87.8)	(12.2)	(100.0)			
総計	5,360	517	5,877	66	2,998	8,941
(%)	(91.2)	(8.8)	(100.0)			
平均	5.24	5.13	5.23	5.64	5.34	5.25
標準偏差	1.33	1.33	1.33	1.69	1.44	1.35

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査