

表 831 心筋梗塞の既往 PTH値(pg/mL)別（腹膜透析患者）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	39	1	40			40
(%)	(97.5)	(2.5)	(100.0)			
20～	44	1	45			45
(%)	(97.8)	(2.2)	(100.0)			
35～	49	5	54	1		55
(%)	(90.7)	(9.3)	(100.0)			
50～	71	5	76	1		77
(%)	(93.4)	(6.6)	(100.0)			
75～	65	4	69			69
(%)	(94.2)	(5.8)	(100.0)			
100～	67	4	71			71
(%)	(94.4)	(5.6)	(100.0)			
125～	39	6	45			45
(%)	(86.7)	(13.3)	(100.0)			
150～	1		1			1
(%)	(100.0)		(100.0)			
151～	37	5	42	1		43
(%)	(88.1)	(11.9)	(100.0)			
180～	30	2	32	2		34
(%)	(93.8)	(6.3)	(100.0)			
210～	25	2	27			27
(%)	(92.6)	(7.4)	(100.0)			
240～	22	1	23			23
(%)	(95.7)	(4.3)	(100.0)			
300～	17	2	19			19
(%)	(89.5)	(10.5)	(100.0)			
350～	8		8			8
(%)	(100.0)		(100.0)			
400～	8	3	11			11
(%)	(72.7)	(27.3)	(100.0)			
500～	10	1	11			11
(%)	(90.9)	(9.1)	(100.0)			
合計	532	42	574	5		579
(%)	(92.7)	(7.3)	(100.0)			
記載なし	100	11	111	29		140
(%)	(90.1)	(9.9)	(100.0)			
総計	632	53	685	34		719
(%)	(92.3)	(7.7)	(100.0)			
平均	129.80	167.67	132.57	131.20		132.56
標準偏差	126.55	163.55	129.82	71.28		129.39

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査