

表 845 脳出血の既往 原疾患別（透析患者全体）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	68,822	4,581	73,403	227	23,340	96,970
(%)	(93.8)	(6.2)	(100.0)			
慢性腎盂腎炎	2,227	133	2,360	5	677	3,042
(%)	(94.4)	(5.6)	(100.0)			
急速進行性糸球体腎炎	1,820	79	1,899	7	588	2,494
(%)	(95.8)	(4.2)	(100.0)			
妊娠腎/妊娠中毒症	1,185	74	1,259	6	349	1,614
(%)	(94.1)	(5.9)	(100.0)			
その他の分類不能の腎炎	971	52	1,023	3	347	1,373
(%)	(94.9)	(5.1)	(100.0)			
多発性嚢胞腎	7,450	1,071	8,521	28	2,457	11,006
(%)	(87.4)	(12.6)	(100.0)			
腎硬化症	20,433	1,604	22,037	121	6,140	28,298
(%)	(92.7)	(7.3)	(100.0)			
悪性高血圧	1,802	146	1,948	6	619	2,573
(%)	(92.5)	(7.5)	(100.0)			
糖尿病性腎症	84,135	4,990	89,125	360	28,596	118,081
(%)	(94.4)	(5.6)	(100.0)			
SLE腎炎	1,637	96	1,733	7	496	2,236
(%)	(94.5)	(5.5)	(100.0)			
アミロイド腎	317	9	326	5	120	451
(%)	(97.2)	(2.8)	(100.0)			
痛風腎	830	49	879	6	228	1,113
(%)	(94.4)	(5.6)	(100.0)			
先天性代謝異常による腎不全	196	9	205	1	66	272
(%)	(95.6)	(4.4)	(100.0)			
腎・尿路結核	145	11	156		50	206
(%)	(92.9)	(7.1)	(100.0)			
腎・尿路結石	408	33	441		140	581
(%)	(92.5)	(7.5)	(100.0)			
腎・尿路腫瘍	629	29	658	1	227	886
(%)	(95.6)	(4.4)	(100.0)			
閉塞性尿路障害	537	26	563	1	159	723
(%)	(95.4)	(4.6)	(100.0)			
骨髄腫	203	8	211	1	64	276
(%)	(96.2)	(3.8)	(100.0)			
腎形成不全	489	21	510		139	649
(%)	(95.9)	(4.1)	(100.0)			
不明	18,802	1,390	20,192	90	7,262	27,544
(%)	(93.1)	(6.9)	(100.0)			
再導入	1,562	81	1,643	2	518	2,163
(%)	(95.1)	(4.9)	(100.0)			
その他	5,322	319	5,641	35	1,881	7,557
(%)	(94.3)	(5.7)	(100.0)			
合計	219,922	14,811	234,733	912	74,463	310,108
(%)	(93.7)	(6.3)	(100.0)			
記載なし						
(%)						
総計	219,922	14,811	234,733	912	74,463	310,108
(%)	(93.7)	(6.3)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査