

表 856 脳出血の既往 身長(cm)別（透析患者全体）

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	293	15	308	2	44	354
(%)	(95.1)	(4.9)	(100.0)			
130～	556	39	595	4	97	696
(%)	(93.4)	(6.6)	(100.0)			
135～	2,187	172	2,359	17	461	2,837
(%)	(92.7)	(7.3)	(100.0)			
140～	8,020	494	8,514	43	1,740	10,297
(%)	(94.2)	(5.8)	(100.0)			
145～	18,352	1,166	19,518	64	4,091	23,673
(%)	(94.0)	(6.0)	(100.0)			
150～	29,433	1,979	31,412	112	6,711	38,235
(%)	(93.7)	(6.3)	(100.0)			
155～	34,405	2,399	36,804	123	7,545	44,472
(%)	(93.5)	(6.5)	(100.0)			
160～	41,269	2,879	44,148	141	9,361	53,650
(%)	(93.5)	(6.5)	(100.0)			
165～	37,161	2,490	39,651	115	8,222	47,988
(%)	(93.7)	(6.3)	(100.0)			
170～	23,971	1,556	25,527	70	5,477	31,074
(%)	(93.9)	(6.1)	(100.0)			
175～	8,796	559	9,355	25	1,939	11,319
(%)	(94.0)	(6.0)	(100.0)			
180～	2,351	108	2,459	9	511	2,979
(%)	(95.6)	(4.4)	(100.0)			
185～	372	12	384	1	80	465
(%)	(96.9)	(3.1)	(100.0)			
190～	61	2	63		14	77
(%)	(96.8)	(3.2)	(100.0)			
合計	207,227	13,870	221,097	726	46,293	268,116
(%)	(93.7)	(6.3)	(100.0)			
記載なし	12,695	941	13,636	186	28,170	41,992
(%)	(93.1)	(6.9)	(100.0)			
総計	219,922	14,811	234,733	912	74,463	310,108
(%)	(93.7)	(6.3)	(100.0)			
平均	160.05	159.90	160.04	158.54	160.09	160.05
標準偏差	9.68	9.48	9.66	10.58	9.58	9.65

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査