

表 864 脳出血の既往 PTH値(pg/mL)別（透析患者全体）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	1,472	125	1,597	1	296	1,894
(%)	(92.2)	(7.8)	(100.0)			
20～	1,833	152	1,985	1	345	2,331
(%)	(92.3)	(7.7)	(100.0)			
35～	1,637	114	1,751	2	290	2,043
(%)	(93.5)	(6.5)	(100.0)			
50～	2,507	141	2,648	3	502	3,153
(%)	(94.7)	(5.3)	(100.0)			
75～	2,146	141	2,287	3	458	2,748
(%)	(93.8)	(6.2)	(100.0)			
100～	1,771	119	1,890	3	405	2,298
(%)	(93.7)	(6.3)	(100.0)			
125～	1,151	79	1,230	1	308	1,539
(%)	(93.6)	(6.4)	(100.0)			
150～	34	4	38		10	48
(%)	(89.5)	(10.5)	(100.0)			
151～	984	60	1,044		219	1,263
(%)	(94.3)	(5.7)	(100.0)			
180～	686	38	724		151	875
(%)	(94.8)	(5.2)	(100.0)			
210～	477	39	516		112	628
(%)	(92.4)	(7.6)	(100.0)			
240～	521	35	556		118	674
(%)	(93.7)	(6.3)	(100.0)			
300～	218	10	228		56	284
(%)	(95.6)	(4.4)	(100.0)			
350～	140	9	149		30	179
(%)	(94.0)	(6.0)	(100.0)			
400～	155	12	167		35	202
(%)	(92.8)	(7.2)	(100.0)			
500～	186	15	201		46	247
(%)	(92.5)	(7.5)	(100.0)			
合計	15,918	1,093	17,011	14	3,381	20,406
(%)	(93.6)	(6.4)	(100.0)			
記載なし	350	44	394	1	262	657
(%)	(88.8)	(11.2)	(100.0)			
総計	16,268	1,137	17,405	15	3,643	21,063
(%)	(93.5)	(6.5)	(100.0)			
平均	106.75	105.48	106.67	70.29	114.89	108.01
標準偏差	110.43	121.66	111.19	35.45	127.26	114.01

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査