

表 871 脳出血の既往 透析前脈拍(拍/分)別 (透析患者全体)

透析前脈拍	既往なし	既往あり	合計	不明	記載なし	総計
40未満	147	7	154		31	185
(%)	(95.5)	(4.5)	(100.0)			
40～	1,855	146	2,001	1	379	2,381
(%)	(92.7)	(7.3)	(100.0)			
50～	15,204	1,101	16,305	57	3,150	19,512
(%)	(93.2)	(6.8)	(100.0)			
60～	47,222	3,377	50,599	158	10,024	60,781
(%)	(93.3)	(6.7)	(100.0)			
70～	57,133	3,746	60,879	203	12,190	73,272
(%)	(93.8)	(6.2)	(100.0)			
80～	36,236	2,263	38,499	154	8,161	46,814
(%)	(94.1)	(5.9)	(100.0)			
90～	16,047	980	17,027	61	3,502	20,590
(%)	(94.2)	(5.8)	(100.0)			
100～	5,079	295	5,374	20	999	6,393
(%)	(94.5)	(5.5)	(100.0)			
110～	1,214	70	1,284	4	240	1,528
(%)	(94.5)	(5.5)	(100.0)			
120～	528	37	565	3	87	655
(%)	(93.5)	(6.5)	(100.0)			
合計	180,665	12,022	192,687	661	38,763	232,111
(%)	(93.8)	(6.2)	(100.0)			
記載なし	39,257	2,789	42,046	251	35,700	77,997
(%)	(93.4)	(6.6)	(100.0)			
総計	219,922	14,811	234,733	912	74,463	310,108
(%)	(93.7)	(6.3)	(100.0)			
平均	74.71	73.89	74.66	75.83	74.83	74.69
標準偏差	12.88	12.76	12.88	12.96	12.63	12.84

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査