

表 875 脳出血の既往 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	52,314	3,611	55,925	176	12,476	68,577
(%)	(93.5)	(6.5)	(100.0)			
慢性腎盂腎炎	1,704	112	1,816	5	344	2,165
(%)	(93.8)	(6.2)	(100.0)			
急速進行性糸球体腎炎	1,474	52	1,526	7	301	1,834
(%)	(96.6)	(3.4)	(100.0)			
妊娠腎/妊娠中毒症	832	56	888	3	173	1,064
(%)	(93.7)	(6.3)	(100.0)			
その他の分類不能の腎炎	722	37	759		190	949
(%)	(95.1)	(4.9)	(100.0)			
多発性嚢胞腎	5,927	882	6,809	23	1,366	8,198
(%)	(87.0)	(13.0)	(100.0)			
腎硬化症	16,660	1,315	17,975	100	3,385	21,460
(%)	(92.7)	(7.3)	(100.0)			
悪性高血圧	1,434	112	1,546	4	331	1,881
(%)	(92.8)	(7.2)	(100.0)			
糖尿病性腎症	69,536	4,115	73,651	294	16,562	90,507
(%)	(94.4)	(5.6)	(100.0)			
SLE腎炎	1,226	69	1,295	4	251	1,550
(%)	(94.7)	(5.3)	(100.0)			
アミロイド腎	249	7	256	5	64	325
(%)	(97.3)	(2.7)	(100.0)			
痛風腎	664	39	703	2	123	828
(%)	(94.5)	(5.5)	(100.0)			
先天性代謝異常による腎不全	142	6	148		29	177
(%)	(95.9)	(4.1)	(100.0)			
腎・尿路結核	122	10	132		26	158
(%)	(92.4)	(7.6)	(100.0)			
腎・尿路結石	324	26	350		80	430
(%)	(92.6)	(7.4)	(100.0)			
腎・尿路腫瘍	528	24	552	1	133	686
(%)	(95.7)	(4.3)	(100.0)			
閉塞性尿路障害	416	20	436	1	80	517
(%)	(95.4)	(4.6)	(100.0)			
骨髄腫	158	6	164	1	25	190
(%)	(96.3)	(3.7)	(100.0)			
腎形成不全	339	14	353		66	419
(%)	(96.0)	(4.0)	(100.0)			
不明	14,973	1,130	16,103	68	3,954	20,125
(%)	(93.0)	(7.0)	(100.0)			
再導入	1,107	60	1,167	2	248	1,417
(%)	(94.9)	(5.1)	(100.0)			
その他	4,235	251	4,486	28	1,000	5,514
(%)	(94.4)	(5.6)	(100.0)			
合計	175,086	11,954	187,040	724	41,207	228,971
(%)	(93.6)	(6.4)	(100.0)			
記載なし						
(%)						
総計	175,086	11,954	187,040	724	41,207	228,971
(%)	(93.6)	(6.4)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査