

表 886 脳出血の既往 身長(cm)別（施設血液透析患者 透析回数週3回）

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	194	11	205	2	30	237
(%)	(94.6)	(5.4)	(100.0)			
130～	467	30	497	2	69	568
(%)	(94.0)	(6.0)	(100.0)			
135～	1,794	135	1,929	16	360	2,305
(%)	(93.0)	(7.0)	(100.0)			
140～	6,577	403	6,980	37	1,266	8,283
(%)	(94.2)	(5.8)	(100.0)			
145～	14,954	932	15,886	50	2,883	18,819
(%)	(94.1)	(5.9)	(100.0)			
150～	23,482	1,627	25,109	96	4,809	30,014
(%)	(93.5)	(6.5)	(100.0)			
155～	27,422	1,958	29,380	98	5,379	34,857
(%)	(93.3)	(6.7)	(100.0)			
160～	33,116	2,318	35,434	123	6,701	42,258
(%)	(93.5)	(6.5)	(100.0)			
165～	29,563	2,041	31,604	83	5,889	37,576
(%)	(93.5)	(6.5)	(100.0)			
170～	18,678	1,254	19,932	53	3,842	23,827
(%)	(93.7)	(6.3)	(100.0)			
175～	6,747	437	7,184	21	1,344	8,549
(%)	(93.9)	(6.1)	(100.0)			
180～	1,811	81	1,892	7	355	2,254
(%)	(95.7)	(4.3)	(100.0)			
185～	287	8	295		56	351
(%)	(97.3)	(2.7)	(100.0)			
190～	47	2	49		8	57
(%)	(95.9)	(4.1)	(100.0)			
合計	165,139	11,237	176,376	588	32,991	209,955
(%)	(93.6)	(6.4)	(100.0)			
記載なし	9,947	717	10,664	136	8,216	19,016
(%)	(93.3)	(6.7)	(100.0)			
総計	175,086	11,954	187,040	724	41,207	228,971
(%)	(93.6)	(6.4)	(100.0)			
平均	159.94	159.88	159.93	158.14	160.03	159.94
標準偏差	9.61	9.38	9.59	10.58	9.54	9.59

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査