

表 887 脳出血の既往 透析後体重(kg)別（施設血液透析患者 透析回数週3回）

透析後体重	既往なし	既往あり	合計	不明	記載なし	総計
20未満	64	5	69		15	84
(%)	(92.8)	(7.2)	(100.0)			
20～	590	74	664	4	134	802
(%)	(88.9)	(11.1)	(100.0)			
30～	14,546	1,374	15,920	57	3,140	19,117
(%)	(91.4)	(8.6)	(100.0)			
40～	45,905	3,655	49,560	202	10,047	59,809
(%)	(92.6)	(7.4)	(100.0)			
50～	55,878	3,719	59,597	206	12,306	72,109
(%)	(93.8)	(6.2)	(100.0)			
60～	34,511	1,989	36,500	125	7,457	44,082
(%)	(94.6)	(5.4)	(100.0)			
70～	13,608	655	14,263	45	2,876	17,184
(%)	(95.4)	(4.6)	(100.0)			
80～	4,858	192	5,050	10	1,021	6,081
(%)	(96.2)	(3.8)	(100.0)			
90～	1,751	74	1,825	5	365	2,195
(%)	(95.9)	(4.1)	(100.0)			
100～	602	19	621	1	133	755
(%)	(96.9)	(3.1)	(100.0)			
110～	246	2	248	1	50	299
(%)	(99.2)	(0.8)	(100.0)			
120～	157	3	160		29	189
(%)	(98.1)	(1.9)	(100.0)			
合計	172,716	11,761	184,477	656	37,573	222,706
(%)	(93.6)	(6.4)	(100.0)			
記載なし	2,370	193	2,563	68	3,634	6,265
(%)	(92.5)	(7.5)	(100.0)			
総計	175,086	11,954	187,040	724	41,207	228,971
(%)	(93.6)	(6.4)	(100.0)			
平均	55.62	52.90	55.44	53.89	55.47	55.45
標準偏差	12.97	11.92	12.92	12.11	12.81	12.90

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査