

表 890 脳出血の既往 Body Mass Index別（施設血液透析患者 透析回数週3回）

Body Mass Index	既往なし	既往あり	合計	不明	記載なし	総計
14未満	943	164	1,107	1	203	1,311
(%)	(85.2)	(14.8)	(100.0)			
14～	5,740	739	6,479	28	1,099	7,606
(%)	(88.6)	(11.4)	(100.0)			
16～	18,994	1,741	20,735	69	3,676	24,480
(%)	(91.6)	(8.4)	(100.0)			
18～	33,950	2,566	36,516	126	6,514	43,156
(%)	(93.0)	(7.0)	(100.0)			
20～	37,767	2,374	40,141	140	7,399	47,680
(%)	(94.1)	(5.9)	(100.0)			
22～	29,223	1,771	30,994	83	5,733	36,810
(%)	(94.3)	(5.7)	(100.0)			
24～	17,575	913	18,488	61	3,364	21,913
(%)	(95.1)	(4.9)	(100.0)			
26～	9,352	465	9,817	33	1,838	11,688
(%)	(95.3)	(4.7)	(100.0)			
28～	4,853	196	5,049	18	924	5,991
(%)	(96.1)	(3.9)	(100.0)			
30～	4,230	162	4,392	12	760	5,164
(%)	(96.3)	(3.7)	(100.0)			
合計	162,627	11,091	173,718	571	31,510	205,799
(%)	(93.6)	(6.4)	(100.0)			
記載なし	12,459	863	13,322	153	9,697	23,172
(%)	(93.5)	(6.5)	(100.0)			
総計	175,086	11,954	187,040	724	41,207	228,971
(%)	(93.6)	(6.4)	(100.0)			
平均	21.52	20.58	21.46	21.29	21.50	21.46
標準偏差	3.66	3.56	3.66	3.61	3.63	3.66

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査