

表 891 脳出血の既往 透析前BUN(mg/dL)別（施設血液透析患者 透析回数週3回 透析歴2年以上）

透析前BUN	既往なし	既往あり	合計	不明	記載なし	総計
20未満	340	50	390	5	84	479
(%)	(87.2)	(12.8)	(100.0)			
20～	1,800	256	2,056	11	413	2,480
(%)	(87.5)	(12.5)	(100.0)			
30～	7,745	822	8,567	33	1,758	10,358
(%)	(90.4)	(9.6)	(100.0)			
40～	20,390	1,852	22,242	84	4,485	26,811
(%)	(91.7)	(8.3)	(100.0)			
50～	32,802	2,473	35,275	102	7,282	42,659
(%)	(93.0)	(7.0)	(100.0)			
60～	32,972	2,316	35,288	101	7,254	42,643
(%)	(93.4)	(6.6)	(100.0)			
70～	22,073	1,418	23,491	68	4,763	28,322
(%)	(94.0)	(6.0)	(100.0)			
80～	10,284	660	10,944	31	2,237	13,212
(%)	(94.0)	(6.0)	(100.0)			
90～	3,725	241	3,966	13	824	4,803
(%)	(93.9)	(6.1)	(100.0)			
100～	1,062	75	1,137	6	205	1,348
(%)	(93.4)	(6.6)	(100.0)			
110～	305	23	328		66	394
(%)	(93.0)	(7.0)	(100.0)			
120～	159	10	169	1	34	204
(%)	(94.1)	(5.9)	(100.0)			
合計	133,657	10,196	143,853	455	29,405	173,713
(%)	(92.9)	(7.1)	(100.0)			
記載なし	1,488	143	1,631	51	2,814	4,496
(%)	(91.2)	(8.8)	(100.0)			
総計	135,145	10,339	145,484	506	32,219	178,209
(%)	(92.9)	(7.1)	(100.0)			
平均	61.12	58.67	60.95	59.16	60.92	60.94
標準偏差	15.59	16.22	15.65	16.88	15.56	15.64

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査