

表 900 脳出血の既往 透析前リン濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

透析前リン濃度	既往なし	既往あり	合計	不明	記載なし	総計
3未満	6,609	774	7,383	31	1,547	8,961
(%)	(89.5)	(10.5)	(100.0)			
3.0～	8,306	739	9,045	34	1,901	10,980
(%)	(91.8)	(8.2)	(100.0)			
3.5～	14,777	1,083	15,860	53	3,105	19,018
(%)	(93.2)	(6.8)	(100.0)			
4.0～	21,949	1,563	23,512	70	4,610	28,192
(%)	(93.4)	(6.6)	(100.0)			
4.5～	26,089	1,773	27,862	90	5,516	33,468
(%)	(93.6)	(6.4)	(100.0)			
5.0～	26,312	1,722	28,034	76	5,699	33,809
(%)	(93.9)	(6.1)	(100.0)			
5.5～	22,701	1,429	24,130	78	4,804	29,012
(%)	(94.1)	(5.9)	(100.0)			
6.0～	3,845	239	4,084	9	835	4,928
(%)	(94.1)	(5.9)	(100.0)			
6.1～	12,932	697	13,629	45	2,831	16,505
(%)	(94.9)	(5.1)	(100.0)			
6.5～	11,268	672	11,940	39	2,398	14,377
(%)	(94.4)	(5.6)	(100.0)			
7.0～	6,926	393	7,319	25	1,516	8,860
(%)	(94.6)	(5.4)	(100.0)			
7.5～	4,204	247	4,451	18	890	5,359
(%)	(94.5)	(5.5)	(100.0)			
8.0～	2,488	148	2,636	6	550	3,192
(%)	(94.4)	(5.6)	(100.0)			
8.5～	1,586	85	1,671	5	347	2,023
(%)	(94.9)	(5.1)	(100.0)			
9.0～	2,419	161	2,580	16	560	3,156
(%)	(93.8)	(6.2)	(100.0)			
合計	172,411	11,725	184,136	595	37,109	221,840
(%)	(93.6)	(6.4)	(100.0)			
記載なし	2,675	229	2,904	129	4,098	7,131
(%)	(92.1)	(7.9)	(100.0)			
総計	175,086	11,954	187,040	724	41,207	228,971
(%)	(93.6)	(6.4)	(100.0)			
平均	5.21	5.01	5.20	5.20	5.21	5.20
標準偏差	1.42	1.48	1.42	1.61	1.44	1.43

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査