

表 902 脳出血の既往 PTH値(pg/mL)別（施設血液透析患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	1,174	100	1,274	1	224	1,499
(%)	(92.2)	(7.8)	(100.0)			
20～	1,489	130	1,619	1	228	1,848
(%)	(92.0)	(8.0)	(100.0)			
35～	1,291	98	1,389	1	223	1,613
(%)	(92.9)	(7.1)	(100.0)			
50～	1,991	118	2,109	3	347	2,459
(%)	(94.4)	(5.6)	(100.0)			
75～	1,709	111	1,820	2	326	2,148
(%)	(93.9)	(6.1)	(100.0)			
100～	1,398	98	1,496	3	288	1,787
(%)	(93.4)	(6.6)	(100.0)			
125～	899	60	959	1	227	1,187
(%)	(93.7)	(6.3)	(100.0)			
150～	27	3	30		7	37
(%)	(90.0)	(10.0)	(100.0)			
151～	738	51	789		145	934
(%)	(93.5)	(6.5)	(100.0)			
180～	526	33	559		107	666
(%)	(94.1)	(5.9)	(100.0)			
210～	351	32	383		89	472
(%)	(91.6)	(8.4)	(100.0)			
240～	411	29	440		86	526
(%)	(93.4)	(6.6)	(100.0)			
300～	157	9	166		39	205
(%)	(94.6)	(5.4)	(100.0)			
350～	102	8	110		25	135
(%)	(92.7)	(7.3)	(100.0)			
400～	119	10	129		23	152
(%)	(92.2)	(7.8)	(100.0)			
500～	130	12	142		24	166
(%)	(91.5)	(8.5)	(100.0)			
合計	12,512	902	13,414	12	2,408	15,834
(%)	(93.3)	(6.7)	(100.0)			
記載なし	218	38	256	1	181	438
(%)	(85.2)	(14.8)	(100.0)			
総計	12,730	940	13,670	13	2,589	16,272
(%)	(93.1)	(6.9)	(100.0)			
平均	104.67	104.40	104.65	71.75	111.67	105.70
標準偏差	109.67	112.02	109.83	37.75	105.97	109.24

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査