

表 903 脳出血の既往 透析前ヘモグロビン濃度(g/dL)別（施設血液透析患者 透析回数週3回）

透析前ヘモグロビン濃度	既往なし	既往あり	合計	不明	記載なし	総計
5未満	217	13	230		31	261
(%)	(94.3)	(5.7)	(100.0)			
5～	290	24	314	1	47	362
(%)	(92.4)	(7.6)	(100.0)			
6～	699	66	765	13	164	942
(%)	(91.4)	(8.6)	(100.0)			
7～	2,747	233	2,980	19	579	3,578
(%)	(92.2)	(7.8)	(100.0)			
8～	9,156	694	9,850	52	1,824	11,726
(%)	(93.0)	(7.0)	(100.0)			
9～	28,831	2,051	30,882	114	6,042	37,038
(%)	(93.4)	(6.6)	(100.0)			
10～	58,735	3,879	62,614	223	12,380	75,217
(%)	(93.8)	(6.2)	(100.0)			
11～	48,098	3,104	51,202	129	10,148	61,479
(%)	(93.9)	(6.1)	(100.0)			
12～	17,413	1,217	18,630	49	3,691	22,370
(%)	(93.5)	(6.5)	(100.0)			
13～	4,291	308	4,599	7	931	5,537
(%)	(93.3)	(6.7)	(100.0)			
14～	1,522	116	1,638	2	293	1,933
(%)	(92.9)	(7.1)	(100.0)			
合計	171,999	11,705	183,704	609	36,130	220,443
(%)	(93.6)	(6.4)	(100.0)			
記載なし	3,087	249	3,336	115	5,077	8,528
(%)	(92.5)	(7.5)	(100.0)			
総計	175,086	11,954	187,040	724	41,207	228,971
(%)	(93.6)	(6.4)	(100.0)			
平均	10.69	10.65	10.69	10.34	10.70	10.69
標準偏差	1.28	1.33	1.28	1.34	1.26	1.28

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査