

表 906 脳出血の既往 Non-HDL-C濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

Non-HDL-C濃度	既往なし	既往あり	合計	不明	記載なし	総計
50未満	2,062	166	2,228	12	476	2,716
(%)	(92.5)	(7.5)	(100.0)			
50～	9,836	685	10,521	54	2,331	12,906
(%)	(93.5)	(6.5)	(100.0)			
70～	21,973	1,531	23,504	87	4,941	28,532
(%)	(93.5)	(6.5)	(100.0)			
90～	25,993	1,720	27,713	96	5,770	33,579
(%)	(93.8)	(6.2)	(100.0)			
110～	20,778	1,380	22,158	69	4,452	26,679
(%)	(93.8)	(6.2)	(100.0)			
130～	12,564	827	13,391	35	2,707	16,133
(%)	(93.8)	(6.2)	(100.0)			
150～	6,196	421	6,617	23	1,403	8,043
(%)	(93.6)	(6.4)	(100.0)			
170～	2,768	194	2,962	9	596	3,567
(%)	(93.5)	(6.5)	(100.0)			
190～	1,042	76	1,118	5	215	1,338
(%)	(93.2)	(6.8)	(100.0)			
210～	613	52	665	5	162	832
(%)	(92.2)	(7.8)	(100.0)			
合計	103,825	7,052	110,877	395	23,053	134,325
(%)	(93.6)	(6.4)	(100.0)			
記載なし	71,261	4,902	76,163	329	18,154	94,646
(%)	(93.6)	(6.4)	(100.0)			
総計	175,086	11,954	187,040	724	41,207	228,971
(%)	(93.6)	(6.4)	(100.0)			
平均	106.55	106.37	106.54	103.04	106.00	106.43
標準偏差	33.16	34.35	33.24	36.79	33.61	33.32

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査