

表 909 脳出血の既往 透析前脈拍(拍/分)別（施設血液透析患者 透析回数週3回）

透析前脈拍	既往なし	既往あり	合計	不明	記載なし	総計
40未満	126	4	130		25	155
(%)	(96.9)	(3.1)	(100.0)			
40～	1,499	128	1,627	1	301	1,929
(%)	(92.1)	(7.9)	(100.0)			
50～	12,405	933	13,338	46	2,389	15,773
(%)	(93.0)	(7.0)	(100.0)			
60～	38,261	2,779	41,040	122	7,579	48,741
(%)	(93.2)	(6.8)	(100.0)			
70～	45,787	3,096	48,883	168	9,133	58,184
(%)	(93.7)	(6.3)	(100.0)			
80～	28,654	1,851	30,505	124	6,094	36,723
(%)	(93.9)	(6.1)	(100.0)			
90～	12,514	776	13,290	42	2,586	15,918
(%)	(94.2)	(5.8)	(100.0)			
100～	3,918	230	4,148	17	739	4,904
(%)	(94.5)	(5.5)	(100.0)			
110～	919	56	975	3	179	1,157
(%)	(94.3)	(5.7)	(100.0)			
120～	404	29	433	3	62	498
(%)	(93.3)	(6.7)	(100.0)			
合計	144,487	9,882	154,369	526	29,087	183,982
(%)	(93.6)	(6.4)	(100.0)			
記載なし	30,599	2,072	32,671	198	12,120	44,989
(%)	(93.7)	(6.3)	(100.0)			
総計	175,086	11,954	187,040	724	41,207	228,971
(%)	(93.6)	(6.4)	(100.0)			
平均	74.51	73.67	74.46	75.72	74.71	74.50
標準偏差	12.83	12.69	12.83	13.08	12.64	12.80

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査