

表 913 脳出血の既往 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	11,576	680	12,256	35	2,785	15,076
(%)	(94.5)	(5.5)	(100.0)			
慢性腎盂腎炎	336	10	346		76	422
(%)	(97.1)	(2.9)	(100.0)			
急速進行性糸球体腎炎	216	13	229		55	284
(%)	(94.3)	(5.7)	(100.0)			
妊娠腎/妊娠中毒症	267	13	280	1	78	359
(%)	(95.4)	(4.6)	(100.0)			
その他の分類不能の腎炎	153	7	160		35	195
(%)	(95.6)	(4.4)	(100.0)			
多発性嚢胞腎	1,062	132	1,194	2	268	1,464
(%)	(88.9)	(11.1)	(100.0)			
腎硬化症	2,213	171	2,384	5	497	2,886
(%)	(92.8)	(7.2)	(100.0)			
悪性高血圧	223	24	247	1	64	312
(%)	(90.3)	(9.7)	(100.0)			
糖尿病性腎症	10,117	572	10,689	33	2,451	13,173
(%)	(94.6)	(5.4)	(100.0)			
SLE腎炎	300	17	317		46	363
(%)	(94.6)	(5.4)	(100.0)			
アミロイド腎	47	2	49		10	59
(%)	(95.9)	(4.1)	(100.0)			
痛風腎	119	6	125	3	26	154
(%)	(95.2)	(4.8)	(100.0)			
先天性代謝異常による腎不全	33	1	34		10	44
(%)	(97.1)	(2.9)	(100.0)			
腎・尿路結核	17	1	18		4	22
(%)	(94.4)	(5.6)	(100.0)			
腎・尿路結石	53	5	58		11	69
(%)	(91.4)	(8.6)	(100.0)			
腎・尿路腫瘍	69	4	73		12	85
(%)	(94.5)	(5.5)	(100.0)			
閉塞性尿路障害	57	5	62		12	74
(%)	(91.9)	(8.1)	(100.0)			
骨髄腫	16		16		3	19
(%)	(100.0)		(100.0)			
腎形成不全	82	4	86		15	101
(%)	(95.3)	(4.7)	(100.0)			
不明	2,568	163	2,731	7	742	3,480
(%)	(94.0)	(6.0)	(100.0)			
再導入	331	18	349		67	416
(%)	(94.8)	(5.2)	(100.0)			
その他	666	38	704	2	183	889
(%)	(94.6)	(5.4)	(100.0)			
合計	30,521	1,886	32,407	89	7,450	39,946
(%)	(94.2)	(5.8)	(100.0)			
記載なし						
(%)						
総計	30,521	1,886	32,407	89	7,450	39,946
(%)	(94.2)	(5.8)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査