

表 924 脳出血の既往 身長(cm)別（血液透析濾過患者 透析回数週3回）

| 身長    | 既往なし    | 既往あり   | 合計      | 不明     | 記載なし   | 総計     |
|-------|---------|--------|---------|--------|--------|--------|
| 130未満 | 33      |        | 33      |        | 5      | 38     |
| (%)   | (100.0) |        | (100.0) |        |        |        |
| 130～  | 46      | 7      | 53      | 1      | 16     | 70     |
| (%)   | (86.8)  | (13.2) | (100.0) |        |        |        |
| 135～  | 222     | 21     | 243     | 1      | 45     | 289    |
| (%)   | (91.4)  | (8.6)  | (100.0) |        |        |        |
| 140～  | 904     | 64     | 968     | 4      | 191    | 1,163  |
| (%)   | (93.4)  | (6.6)  | (100.0) |        |        |        |
| 145～  | 2,307   | 151    | 2,458   | 9      | 528    | 2,995  |
| (%)   | (93.9)  | (6.1)  | (100.0) |        |        |        |
| 150～  | 4,154   | 238    | 4,392   | 8      | 812    | 5,212  |
| (%)   | (94.6)  | (5.4)  | (100.0) |        |        |        |
| 155～  | 4,818   | 306    | 5,124   | 14     | 941    | 6,079  |
| (%)   | (94.0)  | (6.0)  | (100.0) |        |        |        |
| 160～  | 5,652   | 382    | 6,034   | 7      | 1,136  | 7,177  |
| (%)   | (93.7)  | (6.3)  | (100.0) |        |        |        |
| 165～  | 5,359   | 321    | 5,680   | 14     | 1,028  | 6,722  |
| (%)   | (94.3)  | (5.7)  | (100.0) |        |        |        |
| 170～  | 3,833   | 216    | 4,049   | 5      | 731    | 4,785  |
| (%)   | (94.7)  | (5.3)  | (100.0) |        |        |        |
| 175～  | 1,426   | 81     | 1,507   | 2      | 277    | 1,786  |
| (%)   | (94.6)  | (5.4)  | (100.0) |        |        |        |
| 180～  | 402     | 16     | 418     |        | 75     | 493    |
| (%)   | (96.2)  | (3.8)  | (100.0) |        |        |        |
| 185～  | 63      | 3      | 66      |        | 11     | 77     |
| (%)   | (95.5)  | (4.5)  | (100.0) |        |        |        |
| 190～  | 8       |        | 8       |        | 1      | 9      |
| (%)   | (100.0) |        | (100.0) |        |        |        |
| 合計    | 29,227  | 1,806  | 31,033  | 65     | 5,797  | 36,895 |
| (%)   | (94.2)  | (5.8)  | (100.0) |        |        |        |
| 記載なし  | 1,294   | 80     | 1,374   | 24     | 1,653  | 3,051  |
| (%)   | (94.2)  | (5.8)  | (100.0) |        |        |        |
| 総計    | 30,521  | 1,886  | 32,407  | 89     | 7,450  | 39,946 |
| (%)   | (94.2)  | (5.8)  | (100.0) |        |        |        |
| 平均    | 160.82  | 160.29 | 160.79  | 158.08 | 160.48 | 160.74 |
| 標準偏差  | 9.52    | 9.44   | 9.51    | 10.31  | 9.64   | 9.54   |

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査