

表 937 脳出血の既往 透析前補正カルシウム濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

透析前補正 カルシウム濃度	既往なし	既往あり	合計	不明	記載なし	総計
7.0未満	131	6	137		23	160
(%)	(95.6)	(4.4)	(100.0)			
7.0～	211	12	223		36	259
(%)	(94.6)	(5.4)	(100.0)			
7.5～	741	39	780	1	170	951
(%)	(95.0)	(5.0)	(100.0)			
8.0～	2,270	117	2,387	4	477	2,868
(%)	(95.1)	(4.9)	(100.0)			
8.4～	670	24	694	2	122	818
(%)	(96.5)	(3.5)	(100.0)			
8.5～	7,314	423	7,737	14	1,550	9,301
(%)	(94.5)	(5.5)	(100.0)			
9.0～	9,034	539	9,573	24	1,918	11,515
(%)	(94.4)	(5.6)	(100.0)			
9.5～	5,864	401	6,265	11	1,336	7,612
(%)	(93.6)	(6.4)	(100.0)			
10.0～	689	40	729		174	903
(%)	(94.5)	(5.5)	(100.0)			
10.1～	1,877	153	2,030	9	413	2,452
(%)	(92.5)	(7.5)	(100.0)			
10.5～	893	55	948	2	220	1,170
(%)	(94.2)	(5.8)	(100.0)			
11.0～	294	24	318	1	64	383
(%)	(92.5)	(7.5)	(100.0)			
11.5～	99	10	109	2	21	132
(%)	(90.8)	(9.2)	(100.0)			
12.0～	59	5	64		7	71
(%)	(92.2)	(7.8)	(100.0)			
合計	30,146	1,848	31,994	70	6,531	38,595
(%)	(94.2)	(5.8)	(100.0)			
記載なし	375	38	413	19	919	1,351
(%)	(90.8)	(9.2)	(100.0)			
総計	30,521	1,886	32,407	89	7,450	39,946
(%)	(94.2)	(5.8)	(100.0)			
平均	9.18	9.26	9.18	9.36	9.20	9.19
標準偏差	0.75	0.75	0.75	0.76	0.73	0.75

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査