

表 938 脳出血の既往 透析前リン濃度(mg/dL)別（血液透析濾過患者 透析回数週3回）

透析前リン濃度	既往なし	既往あり	合計	不明	記載なし	総計
3未満	876	79	955	1	177	1,133
(%)	(91.7)	(8.3)	(100.0)			
3.0～	1,212	98	1,310	4	260	1,574
(%)	(92.5)	(7.5)	(100.0)			
3.5～	2,220	130	2,350	11	488	2,849
(%)	(94.5)	(5.5)	(100.0)			
4.0～	3,554	244	3,798	9	758	4,565
(%)	(93.6)	(6.4)	(100.0)			
4.5～	4,423	304	4,727	6	979	5,712
(%)	(93.6)	(6.4)	(100.0)			
5.0～	4,784	291	5,075	18	1,041	6,134
(%)	(94.3)	(5.7)	(100.0)			
5.5～	4,157	219	4,376	3	920	5,299
(%)	(95.0)	(5.0)	(100.0)			
6.0～	745	44	789	3	158	950
(%)	(94.4)	(5.6)	(100.0)			
6.1～	2,536	123	2,659	3	582	3,244
(%)	(95.4)	(4.6)	(100.0)			
6.5～	2,182	128	2,310	7	489	2,806
(%)	(94.5)	(5.5)	(100.0)			
7.0～	1,415	77	1,492	3	280	1,775
(%)	(94.8)	(5.2)	(100.0)			
7.5～	870	40	910	1	193	1,104
(%)	(95.6)	(4.4)	(100.0)			
8.0～	514	34	548	1	102	651
(%)	(93.8)	(6.2)	(100.0)			
8.5～	339	13	352		71	423
(%)	(96.3)	(3.7)	(100.0)			
9.0～	534	34	568	1	113	682
(%)	(94.0)	(6.0)	(100.0)			
合計	30,361	1,858	32,219	71	6,611	38,901
(%)	(94.2)	(5.8)	(100.0)			
記載なし	160	28	188	18	839	1,045
(%)	(85.1)	(14.9)	(100.0)			
総計	30,521	1,886	32,407	89	7,450	39,946
(%)	(94.2)	(5.8)	(100.0)			
平均	5.36	5.21	5.35	5.09	5.36	5.36
標準偏差	1.43	1.48	1.43	1.36	1.41	1.43

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査