

表 940 脳出血の既往 PTH値(pg/mL)別（血液透析濾過患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	201	16	217		32	249
(%)	(92.6)	(7.4)	(100.0)			
20～	244	16	260		41	301
(%)	(93.8)	(6.2)	(100.0)			
35～	230	9	239		35	274
(%)	(96.2)	(3.8)	(100.0)			
50～	355	16	371		73	444
(%)	(95.7)	(4.3)	(100.0)			
75～	300	21	321	1	69	391
(%)	(93.5)	(6.5)	(100.0)			
100～	260	19	279		52	331
(%)	(93.2)	(6.8)	(100.0)			
125～	179	14	193		41	234
(%)	(92.7)	(7.3)	(100.0)			
150～	4		4			4
(%)	(100.0)		(100.0)			
151～	179	4	183		32	215
(%)	(97.8)	(2.2)	(100.0)			
180～	113	5	118		19	137
(%)	(95.8)	(4.2)	(100.0)			
210～	83	2	85		13	98
(%)	(97.6)	(2.4)	(100.0)			
240～	72	6	78		10	88
(%)	(92.3)	(7.7)	(100.0)			
300～	36	1	37		5	42
(%)	(97.3)	(2.7)	(100.0)			
350～	27		27		2	29
(%)	(100.0)		(100.0)			
400～	20	1	21		3	24
(%)	(95.2)	(4.8)	(100.0)			
500～	40	2	42		3	45
(%)	(95.2)	(4.8)	(100.0)			
合計	2,343	132	2,475	1	430	2,906
(%)	(94.7)	(5.3)	(100.0)			
記載なし	7	1	8		23	31
(%)	(87.5)	(12.5)	(100.0)			
総計	2,350	133	2,483	1	453	2,937
(%)	(94.6)	(5.4)	(100.0)			
平均	113.22	108.64	112.98	77.00	105.05	111.79
標準偏差	108.07	156.20	111.13		88.76	108.12

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査