

表 941 脳出血の既往 透析前ヘモグロビン濃度(g/dL)別（血液透析濾過患者 透析回数週3回）

透析前ヘモグロビン濃度	既往なし	既往あり	合計	不明	記載なし	総計
5未満	23	1	24		2	26
(%)	(95.8)	(4.2)	(100.0)			
5～	26	1	27	1	3	31
(%)	(96.3)	(3.7)	(100.0)			
6～	92	7	99		15	114
(%)	(92.9)	(7.1)	(100.0)			
7～	363	20	383		62	445
(%)	(94.8)	(5.2)	(100.0)			
8～	1,268	94	1,362	10	265	1,637
(%)	(93.1)	(6.9)	(100.0)			
9～	4,498	307	4,805	19	1,026	5,850
(%)	(93.6)	(6.4)	(100.0)			
10～	10,026	611	10,637	23	2,287	12,947
(%)	(94.3)	(5.7)	(100.0)			
11～	9,054	535	9,589	9	1,979	11,577
(%)	(94.4)	(5.6)	(100.0)			
12～	3,468	184	3,652	7	723	4,382
(%)	(95.0)	(5.0)	(100.0)			
13～	933	61	994	1	190	1,185
(%)	(93.9)	(6.1)	(100.0)			
14～	421	32	453	1	69	523
(%)	(92.9)	(7.1)	(100.0)			
合計	30,172	1,853	32,025	71	6,621	38,717
(%)	(94.2)	(5.8)	(100.0)			
記載なし	349	33	382	18	829	1,229
(%)	(91.4)	(8.6)	(100.0)			
総計	30,521	1,886	32,407	89	7,450	39,946
(%)	(94.2)	(5.8)	(100.0)			
平均	10.85	10.80	10.84	10.28	10.83	10.84
標準偏差	1.27	1.32	1.28	1.42	1.20	1.26

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査