

表 969 脳出血の既往 PTH値(pg/mL)別（腹膜透析患者）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	36	4	40			40
(%)	(90.0)	(10.0)	(100.0)			
20～	45		45			45
(%)	(100.0)		(100.0)			
35～	51	3	54		1	55
(%)	(94.4)	(5.6)	(100.0)			
50～	76		76		1	77
(%)	(100.0)		(100.0)			
75～	66	3	69			69
(%)	(95.7)	(4.3)	(100.0)			
100～	69	2	71			71
(%)	(97.2)	(2.8)	(100.0)			
125～	42	3	45			45
(%)	(93.3)	(6.7)	(100.0)			
150～	1		1			1
(%)	(100.0)		(100.0)			
151～	39	3	42		1	43
(%)	(92.9)	(7.1)	(100.0)			
180～	32		32		2	34
(%)	(100.0)		(100.0)			
210～	25	2	27			27
(%)	(92.6)	(7.4)	(100.0)			
240～	23		23			23
(%)	(100.0)		(100.0)			
300～	19		19			19
(%)	(100.0)		(100.0)			
350～	8		8			8
(%)	(100.0)		(100.0)			
400～	10	1	11			11
(%)	(90.9)	(9.1)	(100.0)			
500～	10	1	11			11
(%)	(90.9)	(9.1)	(100.0)			
合計	552	22	574		5	579
(%)	(96.2)	(3.8)	(100.0)			
記載なし	109	1	110		30	140
(%)	(99.1)	(0.9)	(100.0)			
総計	661	23	684		35	719
(%)	(96.6)	(3.4)	(100.0)			
平均	131.27	165.09	132.57		131.20	132.56
標準偏差	122.69	252.47	129.82		71.28	129.39

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査