

表 993 脳梗塞の既往 身長(cm)別（透析患者全体）

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	257	57	314	2	38	354
(%)	(81.8)	(18.2)	(100.0)			
130～	482	116	598	4	94	696
(%)	(80.6)	(19.4)	(100.0)			
135～	1,911	487	2,398	20	419	2,837
(%)	(79.7)	(20.3)	(100.0)			
140～	6,957	1,727	8,684	57	1,556	10,297
(%)	(80.1)	(19.9)	(100.0)			
145～	16,158	3,681	19,839	84	3,750	23,673
(%)	(81.4)	(18.6)	(100.0)			
150～	26,093	5,813	31,906	133	6,196	38,235
(%)	(81.8)	(18.2)	(100.0)			
155～	30,259	7,095	37,354	132	6,986	44,472
(%)	(81.0)	(19.0)	(100.0)			
160～	36,051	8,952	45,003	155	8,492	53,650
(%)	(80.1)	(19.9)	(100.0)			
165～	32,762	7,503	40,265	126	7,597	47,988
(%)	(81.4)	(18.6)	(100.0)			
170～	21,659	4,240	25,899	72	5,103	31,074
(%)	(83.6)	(16.4)	(100.0)			
175～	8,122	1,336	9,458	21	1,840	11,319
(%)	(85.9)	(14.1)	(100.0)			
180～	2,197	291	2,488	7	484	2,979
(%)	(88.3)	(11.7)	(100.0)			
185～	351	33	384	1	80	465
(%)	(91.4)	(8.6)	(100.0)			
190～	56	7	63		14	77
(%)	(88.9)	(11.1)	(100.0)			
合計	183,315	41,338	224,653	814	42,649	268,116
(%)	(81.6)	(18.4)	(100.0)			
記載なし	11,326	2,794	14,120	166	27,706	41,992
(%)	(80.2)	(19.8)	(100.0)			
総計	194,641	44,132	238,773	980	70,355	310,108
(%)	(81.5)	(18.5)	(100.0)			
平均	160.14	159.52	160.03	157.86	160.17	160.05
標準偏差	9.72	9.36	9.66	10.38	9.60	9.65

数値下のカッコ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査