

表 1002 脳梗塞の既往 透析前ヘモグロビン濃度(g/dL)別 (透析患者全体)

透析前ヘモグロビン濃度	既往なし	既往あり	合計	不明	記載なし	総計
5未満	220	53	273		30	303
(%)	(80.6)	(19.4)	(100.0)			
5～	298	71	369	2	58	429
(%)	(80.8)	(19.2)	(100.0)			
6～	756	212	968	11	181	1,160
(%)	(78.1)	(21.9)	(100.0)			
7～	2,943	794	3,737	17	681	4,435
(%)	(78.8)	(21.2)	(100.0)			
8～	9,669	2,627	12,296	74	2,184	14,554
(%)	(78.6)	(21.4)	(100.0)			
9～	31,059	7,506	38,565	158	7,266	45,989
(%)	(80.5)	(19.5)	(100.0)			
10～	64,254	14,186	78,440	311	15,161	93,912
(%)	(81.9)	(18.1)	(100.0)			
11～	54,107	11,213	65,320	221	12,632	78,173
(%)	(82.8)	(17.2)	(100.0)			
12～	19,916	4,294	24,210	76	4,568	28,854
(%)	(82.3)	(17.7)	(100.0)			
13～	5,047	1,082	6,129	14	1,165	7,308
(%)	(82.3)	(17.7)	(100.0)			
14～	1,876	408	2,284	3	386	2,673
(%)	(82.1)	(17.9)	(100.0)			
合計	190,145	42,446	232,591	887	44,312	277,790
(%)	(81.8)	(18.2)	(100.0)			
記載なし	4,496	1,686	6,182	93	26,043	32,318
(%)	(72.7)	(27.3)	(100.0)			
総計	194,641	44,132	238,773	980	70,355	310,108
(%)	(81.5)	(18.5)	(100.0)			
平均	10.72	10.64	10.71	10.47	10.72	10.71
標準偏差	1.28	1.32	1.29	1.29	1.25	1.28

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査