

表 1008 脳梗塞の既往 透析前脈拍(拍/分)別 (透析患者全体)

透析前脈拍	既往なし	既往あり	合計	不明	記載なし	総計
40未満	111	42	153		32	185
(%)	(72.5)	(27.5)	(100.0)			
40～	1,551	488	2,039	3	339	2,381
(%)	(76.1)	(23.9)	(100.0)			
50～	13,137	3,416	16,553	66	2,893	19,512
(%)	(79.4)	(20.6)	(100.0)			
60～	41,779	9,548	51,327	192	9,262	60,781
(%)	(81.4)	(18.6)	(100.0)			
70～	50,837	10,871	61,708	238	11,326	73,272
(%)	(82.4)	(17.6)	(100.0)			
80～	32,159	6,834	38,993	172	7,649	46,814
(%)	(82.5)	(17.5)	(100.0)			
90～	14,236	3,011	17,247	74	3,269	20,590
(%)	(82.5)	(17.5)	(100.0)			
100～	4,464	984	5,448	20	925	6,393
(%)	(81.9)	(18.1)	(100.0)			
110～	1,078	221	1,299	7	222	1,528
(%)	(83.0)	(17.0)	(100.0)			
120～	466	106	572	2	81	655
(%)	(81.5)	(18.5)	(100.0)			
合計	159,818	35,521	195,339	774	35,998	232,111
(%)	(81.8)	(18.2)	(100.0)			
記載なし	34,823	8,611	43,434	206	34,357	77,997
(%)	(80.2)	(19.8)	(100.0)			
総計	194,641	44,132	238,773	980	70,355	310,108
(%)	(81.5)	(18.5)	(100.0)			
平均	74.77	74.08	74.65	75.48	74.89	74.69
標準偏差	12.81	13.16	12.88	12.82	12.61	12.84

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査