

表 1012 脳梗塞の既往 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	既往なし	既往あり	合計	不明	記載なし	総計
慢性糸球体腎炎	47,949	8,560	56,509	217	11,851	68,577
(%)	(84.9)	(15.1)	(100.0)			
慢性腎盂腎炎	1,591	237	1,828	6	331	2,165
(%)	(87.0)	(13.0)	(100.0)			
急速進行性糸球体腎炎	1,355	182	1,537	7	290	1,834
(%)	(88.2)	(11.8)	(100.0)			
妊娠腎/妊娠中毒症	769	128	897	2	165	1,064
(%)	(85.7)	(14.3)	(100.0)			
その他の分類不能の腎炎	689	75	764		185	949
(%)	(90.2)	(9.8)	(100.0)			
多発性嚢胞腎	5,828	979	6,807	27	1,364	8,198
(%)	(85.6)	(14.4)	(100.0)			
腎硬化症	14,535	3,665	18,200	101	3,159	21,460
(%)	(79.9)	(20.1)	(100.0)			
悪性高血圧	1,305	263	1,568	2	311	1,881
(%)	(83.2)	(16.8)	(100.0)			
糖尿病性腎症	58,126	17,114	75,240	317	14,950	90,507
(%)	(77.3)	(22.7)	(100.0)			
SLE腎炎	1,130	173	1,303	6	241	1,550
(%)	(86.7)	(13.3)	(100.0)			
アミロイド腎	232	29	261	2	62	325
(%)	(88.9)	(11.1)	(100.0)			
痛風腎	603	100	703	2	123	828
(%)	(85.8)	(14.2)	(100.0)			
先天性代謝異常による腎不全	130	20	150		27	177
(%)	(86.7)	(13.3)	(100.0)			
腎・尿路結核	113	18	131		27	158
(%)	(86.3)	(13.7)	(100.0)			
腎・尿路結石	282	76	358		72	430
(%)	(78.8)	(21.2)	(100.0)			
腎・尿路腫瘍	477	78	555	2	129	686
(%)	(85.9)	(14.1)	(100.0)			
閉塞性尿路障害	400	43	443	1	73	517
(%)	(90.3)	(9.7)	(100.0)			
骨髄腫	152	14	166		24	190
(%)	(91.6)	(8.4)	(100.0)			
腎形成不全	334	20	354	1	64	419
(%)	(94.4)	(5.6)	(100.0)			
不明	13,438	2,940	16,378	75	3,672	20,125
(%)	(82.0)	(18.0)	(100.0)			
再導入	1,047	126	1,173	3	241	1,417
(%)	(89.3)	(10.7)	(100.0)			
その他	3,880	652	4,532	29	953	5,514
(%)	(85.6)	(14.4)	(100.0)			
合計	154,365	35,492	189,857	800	38,314	228,971
(%)	(81.3)	(18.7)	(100.0)			
記載なし						
(%)						
総計	154,365	35,492	189,857	800	38,314	228,971
(%)	(81.3)	(18.7)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査