

表 1022 脳梗塞の既往 身長(cm)別（施設血液透析患者 透析回数週3回）

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	163	46	209	2	26	237
(%)	(78.0)	(22.0)	(100.0)			
130～	406	93	499	1	68	568
(%)	(81.4)	(18.6)	(100.0)			
135～	1,554	406	1,960	19	326	2,305
(%)	(79.3)	(20.7)	(100.0)			
140～	5,683	1,408	7,091	51	1,141	8,283
(%)	(80.1)	(19.9)	(100.0)			
145～	13,114	2,997	16,111	70	2,638	18,819
(%)	(81.4)	(18.6)	(100.0)			
150～	20,768	4,691	25,459	115	4,440	30,014
(%)	(81.6)	(18.4)	(100.0)			
155～	23,993	5,780	29,773	112	4,972	34,857
(%)	(80.6)	(19.4)	(100.0)			
160～	28,786	7,249	36,035	132	6,091	42,258
(%)	(79.9)	(20.1)	(100.0)			
165～	25,961	6,087	32,048	98	5,430	37,576
(%)	(81.0)	(19.0)	(100.0)			
170～	16,834	3,361	20,195	58	3,574	23,827
(%)	(83.4)	(16.6)	(100.0)			
175～	6,214	1,040	7,254	17	1,278	8,549
(%)	(85.7)	(14.3)	(100.0)			
180～	1,702	210	1,912	6	336	2,254
(%)	(89.0)	(11.0)	(100.0)			
185～	271	25	296		55	351
(%)	(91.6)	(8.4)	(100.0)			
190～	45	4	49		8	57
(%)	(91.8)	(8.2)	(100.0)			
合計	145,494	33,397	178,891	681	30,383	209,955
(%)	(81.3)	(18.7)	(100.0)			
記載なし	8,871	2,095	10,966	119	7,931	19,016
(%)	(80.9)	(19.1)	(100.0)			
総計	154,365	35,492	189,857	800	38,314	228,971
(%)	(81.3)	(18.7)	(100.0)			
平均	160.04	159.44	159.93	157.54	160.10	159.94
標準偏差	9.65	9.32	9.59	10.34	9.55	9.59

数値下のカッコ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2014年調査