

表 1023 脳梗塞の既往 透析後体重(kg)別（施設血液透析患者 透析回数週3回）

透析後体重	既往なし	既往あり	合計	不明	記載なし	総計
20未満	49	24	73		11	84
(%)	(67.1)	(32.9)	(100.0)			
20～	493	183	676	4	122	802
(%)	(72.9)	(27.1)	(100.0)			
30～	12,679	3,479	16,158	65	2,894	19,117
(%)	(78.5)	(21.5)	(100.0)			
40～	40,129	10,173	50,302	212	9,295	59,809
(%)	(79.8)	(20.2)	(100.0)			
50～	48,774	11,771	60,545	251	11,313	72,109
(%)	(80.6)	(19.4)	(100.0)			
60～	30,665	6,307	36,972	140	6,970	44,082
(%)	(82.9)	(17.1)	(100.0)			
70～	12,305	2,108	14,413	42	2,729	17,184
(%)	(85.4)	(14.6)	(100.0)			
80～	4,493	607	5,100	14	967	6,081
(%)	(88.1)	(11.9)	(100.0)			
90～	1,658	172	1,830	8	357	2,195
(%)	(90.6)	(9.4)	(100.0)			
100～	570	55	625	1	129	755
(%)	(91.2)	(8.8)	(100.0)			
110～	231	18	249	1	49	299
(%)	(92.8)	(7.2)	(100.0)			
120～	153	8	161		28	189
(%)	(95.0)	(5.0)	(100.0)			
合計	152,199	34,905	187,104	738	34,864	222,706
(%)	(81.3)	(18.7)	(100.0)			
記載なし	2,166	587	2,753	62	3,450	6,265
(%)	(78.7)	(21.3)	(100.0)			
総計	154,365	35,492	189,857	800	38,314	228,971
(%)	(81.3)	(18.7)	(100.0)			
平均	55.82	53.68	55.42	54.29	55.60	55.45
標準偏差	13.11	11.79	12.90	12.00	12.91	12.90

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査