

表 1026 脳梗塞の既往 Body Mass Index別（施設血液透析患者 透析回数週3回）

Body Mass Index	既往なし	既往あり	合計	不明	記載なし	総計
14未満	775	342	1,117	2	192	1,311
(%)	(69.4)	(30.6)	(100.0)			
14～	4,919	1,686	6,605	23	978	7,606
(%)	(74.5)	(25.5)	(100.0)			
16～	16,579	4,434	21,013	76	3,391	24,480
(%)	(78.9)	(21.1)	(100.0)			
18～	29,866	7,177	37,043	126	5,987	43,156
(%)	(80.6)	(19.4)	(100.0)			
20～	33,087	7,656	40,743	164	6,773	47,680
(%)	(81.2)	(18.8)	(100.0)			
22～	25,889	5,520	31,409	119	5,282	36,810
(%)	(82.4)	(17.6)	(100.0)			
24～	15,505	3,207	18,712	76	3,125	21,913
(%)	(82.9)	(17.1)	(100.0)			
26～	8,352	1,579	9,931	44	1,713	11,688
(%)	(84.1)	(15.9)	(100.0)			
28～	4,356	748	5,104	19	868	5,991
(%)	(85.3)	(14.7)	(100.0)			
30～	3,870	543	4,413	16	735	5,164
(%)	(87.7)	(12.3)	(100.0)			
合計	143,198	32,892	176,090	665	29,044	205,799
(%)	(81.3)	(18.7)	(100.0)			
記載なし	11,167	2,600	13,767	135	9,270	23,172
(%)	(81.1)	(18.9)	(100.0)			
総計	154,365	35,492	189,857	800	38,314	228,971
(%)	(81.3)	(18.7)	(100.0)			
平均	21.55	21.00	21.45	21.61	21.53	21.46
標準偏差	3.67	3.56	3.66	3.56	3.66	3.66

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査