

表 1038 脳梗塞の既往 PTH値(pg/mL)別（施設血液透析患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	1,031	266	1,297	3	199	1,499
(%)	(79.5)	(20.5)	(100.0)			
20～	1,293	339	1,632	1	215	1,848
(%)	(79.2)	(20.8)	(100.0)			
35～	1,135	270	1,405	2	206	1,613
(%)	(80.8)	(19.2)	(100.0)			
50～	1,714	419	2,133	3	323	2,459
(%)	(80.4)	(19.6)	(100.0)			
75～	1,485	372	1,857	1	290	2,148
(%)	(80.0)	(20.0)	(100.0)			
100～	1,215	305	1,520	1	266	1,787
(%)	(79.9)	(20.1)	(100.0)			
125～	780	197	977		210	1,187
(%)	(79.8)	(20.2)	(100.0)			
150～	26	3	29		8	37
(%)	(89.7)	(10.3)	(100.0)			
151～	656	147	803		131	934
(%)	(81.7)	(18.3)	(100.0)			
180～	462	110	572		94	666
(%)	(80.8)	(19.2)	(100.0)			
210～	323	72	395		77	472
(%)	(81.8)	(18.2)	(100.0)			
240～	377	71	448		78	526
(%)	(84.2)	(15.8)	(100.0)			
300～	142	27	169		36	205
(%)	(84.0)	(16.0)	(100.0)			
350～	94	19	113		22	135
(%)	(83.2)	(16.8)	(100.0)			
400～	119	12	131		21	152
(%)	(90.8)	(9.2)	(100.0)			
500～	119	32	151	1	14	166
(%)	(78.8)	(21.2)	(100.0)			
合計	10,971	2,661	13,632	12	2,190	15,834
(%)	(80.5)	(19.5)	(100.0)			
記載なし	198	60	258	1	179	438
(%)	(76.7)	(23.3)	(100.0)			
総計	11,169	2,721	13,890	13	2,369	16,272
(%)	(80.4)	(19.6)	(100.0)			
平均	106.51	100.22	105.28	99.83	108.31	105.70
標準偏差	113.78	102.13	111.63	188.05	92.39	109.24

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査