

表 1039 脳梗塞の既往 透析前ヘモグロビン濃度(g/dL)別（施設血液透析患者 透析回数週3回）

透析前ヘモグロビン濃度	既往なし	既往あり	合計	不明	記載なし	総計
5未満	191	44	235		26	261
(%)	(81.3)	(18.7)	(100.0)			
5～	254	64	318	1	43	362
(%)	(79.9)	(20.1)	(100.0)			
6～	608	177	785	11	146	942
(%)	(77.5)	(22.5)	(100.0)			
7～	2,386	656	3,042	16	520	3,578
(%)	(78.4)	(21.6)	(100.0)			
8～	7,878	2,145	10,023	58	1,645	11,726
(%)	(78.6)	(21.4)	(100.0)			
9～	25,158	6,205	31,363	121	5,554	37,038
(%)	(80.2)	(19.8)	(100.0)			
10～	51,767	11,689	63,456	266	11,495	75,217
(%)	(81.6)	(18.4)	(100.0)			
11～	42,655	9,169	51,824	186	9,469	61,479
(%)	(82.3)	(17.7)	(100.0)			
12～	15,414	3,480	18,894	64	3,412	22,370
(%)	(81.6)	(18.4)	(100.0)			
13～	3,844	824	4,668	11	858	5,537
(%)	(82.3)	(17.7)	(100.0)			
14～	1,347	313	1,660	2	271	1,933
(%)	(81.1)	(18.9)	(100.0)			
合計	151,502	34,766	186,268	736	33,439	220,443
(%)	(81.3)	(18.7)	(100.0)			
記載なし	2,863	726	3,589	64	4,875	8,528
(%)	(79.8)	(20.2)	(100.0)			
総計	154,365	35,492	189,857	800	38,314	228,971
(%)	(81.3)	(18.7)	(100.0)			
平均	10.70	10.63	10.69	10.49	10.71	10.69
標準偏差	1.27	1.31	1.28	1.29	1.25	1.28

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査