

表 1042 脳梗塞の既往 Non-HDL-C濃度(mg/dL)別（施設血液透析患者 透析回数週3回）

Non-HDL-C濃度	既往なし	既往あり	合計	不明	記載なし	総計
50未満	1,822	434	2,256	12	448	2,716
(%)	(80.8)	(19.2)	(100.0)			
50～	8,513	2,185	10,698	59	2,149	12,906
(%)	(79.6)	(20.4)	(100.0)			
70～	19,232	4,659	23,891	109	4,532	28,532
(%)	(80.5)	(19.5)	(100.0)			
90～	22,765	5,364	28,129	127	5,323	33,579
(%)	(80.9)	(19.1)	(100.0)			
110～	18,335	4,138	22,473	85	4,121	26,679
(%)	(81.6)	(18.4)	(100.0)			
130～	11,104	2,468	13,572	44	2,517	16,133
(%)	(81.8)	(18.2)	(100.0)			
150～	5,554	1,157	6,711	32	1,300	8,043
(%)	(82.8)	(17.2)	(100.0)			
170～	2,493	505	2,998	10	559	3,567
(%)	(83.2)	(16.8)	(100.0)			
190～	927	207	1,134	7	197	1,338
(%)	(81.7)	(18.3)	(100.0)			
210～	543	128	671	5	156	832
(%)	(80.9)	(19.1)	(100.0)			
合計	91,288	21,245	112,533	490	21,302	134,325
(%)	(81.1)	(18.9)	(100.0)			
記載なし	63,077	14,247	77,324	310	17,012	94,646
(%)	(81.6)	(18.4)	(100.0)			
総計	154,365	35,492	189,857	800	38,314	228,971
(%)	(81.3)	(18.7)	(100.0)			
平均	106.81	105.13	106.50	104.08	106.17	106.43
標準偏差	33.25	33.13	33.23	35.27	33.69	33.32

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査