

表 1045 脳梗塞の既往 透析前脈拍(拍/分)別（施設血液透析患者 透析回数週3回）

透析前脈拍	既往なし	既往あり	合計	不明	記載なし	総計
40未満	96	34	130		25	155
(%)	(73.8)	(26.2)	(100.0)			
40～	1,254	401	1,655	3	271	1,929
(%)	(75.8)	(24.2)	(100.0)			
50～	10,671	2,857	13,528	56	2,189	15,773
(%)	(78.9)	(21.1)	(100.0)			
60～	33,690	7,906	41,596	156	6,989	48,741
(%)	(81.0)	(19.0)	(100.0)			
70～	40,607	8,912	49,519	201	8,464	58,184
(%)	(82.0)	(18.0)	(100.0)			
80～	25,291	5,603	30,894	139	5,690	36,723
(%)	(81.9)	(18.1)	(100.0)			
90～	11,016	2,434	13,450	57	2,411	15,918
(%)	(81.9)	(18.1)	(100.0)			
100～	3,431	775	4,206	15	683	4,904
(%)	(81.6)	(18.4)	(100.0)			
110～	814	168	982	6	169	1,157
(%)	(82.9)	(17.1)	(100.0)			
120～	359	80	439	2	57	498
(%)	(81.8)	(18.2)	(100.0)			
合計	127,229	29,170	156,399	635	26,948	183,982
(%)	(81.3)	(18.7)	(100.0)			
記載なし	27,136	6,322	33,458	165	11,366	44,989
(%)	(81.1)	(18.9)	(100.0)			
総計	154,365	35,492	189,857	800	38,314	228,971
(%)	(81.3)	(18.7)	(100.0)			
平均	74.57	73.92	74.45	75.23	74.77	74.50
標準偏差	12.76	13.09	12.83	12.88	12.63	12.80

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査