

表 1059 脳梗塞の既往 身長(cm)別（血液透析濾過患者 透析回数週3回）

身長	既往なし	既往あり	合計	不明	記載なし	総計
130未満	30	5	35		3	38
(%)	(85.7)	(14.3)	(100.0)			
130～	41	11	52	1	17	70
(%)	(78.8)	(21.2)	(100.0)			
135～	201	46	247	1	41	289
(%)	(81.4)	(18.6)	(100.0)			
140～	798	192	990	4	169	1,163
(%)	(80.6)	(19.4)	(100.0)			
145～	2,061	439	2,500	9	486	2,995
(%)	(82.4)	(17.6)	(100.0)			
150～	3,734	713	4,447	9	756	5,212
(%)	(84.0)	(16.0)	(100.0)			
155～	4,328	885	5,213	11	855	6,079
(%)	(83.0)	(17.0)	(100.0)			
160～	5,010	1,133	6,143	11	1,023	7,177
(%)	(81.6)	(18.4)	(100.0)			
165～	4,795	960	5,755	11	956	6,722
(%)	(83.3)	(16.7)	(100.0)			
170～	3,492	595	4,087	5	693	4,785
(%)	(85.4)	(14.6)	(100.0)			
175～	1,323	198	1,521	2	263	1,786
(%)	(87.0)	(13.0)	(100.0)			
180～	364	59	423		70	493
(%)	(86.1)	(13.9)	(100.0)			
185～	59	6	65		12	77
(%)	(90.8)	(9.2)	(100.0)			
190～	7	1	8		1	9
(%)	(87.5)	(12.5)	(100.0)			
合計	26,243	5,243	31,486	64	5,345	36,895
(%)	(83.3)	(16.7)	(100.0)			
記載なし	1,159	247	1,406	20	1,625	3,051
(%)	(82.4)	(17.6)	(100.0)			
総計	27,402	5,490	32,892	84	6,970	39,946
(%)	(83.3)	(16.7)	(100.0)			
平均	160.88	160.22	160.77	157.77	160.58	160.74
標準偏差	9.54	9.31	9.51	10.21	9.69	9.54

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査