

表 1060 脳梗塞の既往 透析後体重(kg)別（血液透析濾過患者 透析回数週3回）

透析後体重	既往なし	既往あり	合計	不明	記載なし	総計
20未満	7	2	9		3	12
(%)	(77.8)	(22.2)	(100.0)			
20～	52	10	62	1	13	76
(%)	(83.9)	(16.1)	(100.0)			
30～	1,800	439	2,239	8	435	2,682
(%)	(80.4)	(19.6)	(100.0)			
40～	6,920	1,444	8,364	29	1,612	10,005
(%)	(82.7)	(17.3)	(100.0)			
50～	8,671	1,814	10,485	19	1,986	12,490
(%)	(82.7)	(17.3)	(100.0)			
60～	5,868	1,143	7,011	14	1,340	8,365
(%)	(83.7)	(16.3)	(100.0)			
70～	2,489	399	2,888	2	533	3,423
(%)	(86.2)	(13.8)	(100.0)			
80～	940	138	1,078		195	1,273
(%)	(87.2)	(12.8)	(100.0)			
90～	340	40	380		75	455
(%)	(89.5)	(10.5)	(100.0)			
100～	99	14	113		29	142
(%)	(87.6)	(12.4)	(100.0)			
110～	51	1	52		9	61
(%)	(98.1)	(1.9)	(100.0)			
120～	30	2	32		5	37
(%)	(93.8)	(6.3)	(100.0)			
合計	27,267	5,446	32,713	73	6,235	39,021
(%)	(83.4)	(16.6)	(100.0)			
記載なし	135	44	179	11	735	925
(%)	(75.4)	(24.6)	(100.0)			
総計	27,402	5,490	32,892	84	6,970	39,946
(%)	(83.3)	(16.7)	(100.0)			
平均	56.92	55.39	56.66	51.24	56.49	56.63
標準偏差	13.19	12.15	13.03	10.03	13.05	13.03

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査