

表 1064 脳梗塞の既往 透析前BUN(mg/dL)別（血液透析濾過患者 透析回数週3回 透析歴2年以上）

透析前BUN	既往なし	既往あり	合計	不明	記載なし	総計
20未満	24	7	31		8	39
(%)	(77.4)	(22.6)	(100.0)			
20～	194	74	268		41	309
(%)	(72.4)	(27.6)	(100.0)			
30～	1,068	340	1,408	3	242	1,653
(%)	(75.9)	(24.1)	(100.0)			
40～	3,218	832	4,050	15	720	4,785
(%)	(79.5)	(20.5)	(100.0)			
50～	5,705	1,281	6,986	9	1,323	8,318
(%)	(81.7)	(18.3)	(100.0)			
60～	6,237	1,153	7,390	14	1,400	8,804
(%)	(84.4)	(15.6)	(100.0)			
70～	4,283	741	5,024	9	1,053	6,086
(%)	(85.3)	(14.7)	(100.0)			
80～	2,041	314	2,355	4	472	2,831
(%)	(86.7)	(13.3)	(100.0)			
90～	754	116	870	1	180	1,051
(%)	(86.7)	(13.3)	(100.0)			
100～	206	27	233	1	45	279
(%)	(88.4)	(11.6)	(100.0)			
110～	48	8	56		18	74
(%)	(85.7)	(14.3)	(100.0)			
120～	10	6	16		4	20
(%)	(62.5)	(37.5)	(100.0)			
合計	23,788	4,899	28,687	56	5,506	34,249
(%)	(82.9)	(17.1)	(100.0)			
記載なし	90	38	128	10	584	722
(%)	(70.3)	(29.7)	(100.0)			
総計	23,878	4,937	28,815	66	6,090	34,971
(%)	(82.9)	(17.1)	(100.0)			
平均	62.61	59.66	62.11	59.63	62.91	62.23
標準偏差	14.98	15.30	15.08	15.28	15.10	15.08

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査