

表 1075 脳梗塞の既往 PTH値(pg/mL)別（血液透析濾過患者 透析回数週3回）PTH測定法:whole-PTH

PTH値	既往なし	既往あり	合計	不明	記載なし	総計
20未満	182	35	217		32	249
(%)	(83.9)	(16.1)	(100.0)			
20～	222	42	264		37	301
(%)	(84.1)	(15.9)	(100.0)			
35～	200	39	239		35	274
(%)	(83.7)	(16.3)	(100.0)			
50～	319	55	374		70	444
(%)	(85.3)	(14.7)	(100.0)			
75～	276	58	334	1	56	391
(%)	(82.6)	(17.4)	(100.0)			
100～	235	44	279		52	331
(%)	(84.2)	(15.8)	(100.0)			
125～	152	50	202		32	234
(%)	(75.2)	(24.8)	(100.0)			
150～	4		4			4
(%)	(100.0)		(100.0)			
151～	157	33	190		25	215
(%)	(82.6)	(17.4)	(100.0)			
180～	103	19	122		15	137
(%)	(84.4)	(15.6)	(100.0)			
210～	72	16	88		10	98
(%)	(81.8)	(18.2)	(100.0)			
240～	61	18	79		9	88
(%)	(77.2)	(22.8)	(100.0)			
300～	33	4	37		5	42
(%)	(89.2)	(10.8)	(100.0)			
350～	25	2	27		2	29
(%)	(92.6)	(7.4)	(100.0)			
400～	16	5	21		3	24
(%)	(76.2)	(23.8)	(100.0)			
500～	38	5	43		2	45
(%)	(88.4)	(11.6)	(100.0)			
合計	2,095	425	2,520	1	385	2,906
(%)	(83.1)	(16.9)	(100.0)			
記載なし	8		8		23	31
(%)	(100.0)		(100.0)			
総計	2,103	425	2,528	1	408	2,937
(%)	(83.2)	(16.8)	(100.0)			
平均	113.30	113.73	113.37	77.00	101.51	111.79
標準偏差	113.02	98.31	110.66		89.29	108.12

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査