

表 1082 脳梗塞の既往 透析前脈拍(拍/分)別（血液透析濾過患者 透析回数週3回）

透析前脈拍	既往なし	既往あり	合計	不明	記載なし	総計
40未満	8	6	14		3	17
(%)	(57.1)	(42.9)	(100.0)			
40～	208	53	261		34	295
(%)	(79.7)	(20.3)	(100.0)			
50～	1,825	388	2,213	7	412	2,632
(%)	(82.5)	(17.5)	(100.0)			
60～	5,926	1,199	7,125	18	1,347	8,490
(%)	(83.2)	(16.8)	(100.0)			
70～	7,423	1,401	8,824	19	1,702	10,545
(%)	(84.1)	(15.9)	(100.0)			
80～	4,935	908	5,843	18	1,202	7,063
(%)	(84.5)	(15.5)	(100.0)			
90～	2,314	426	2,740	8	545	3,293
(%)	(84.5)	(15.5)	(100.0)			
100～	721	151	872	2	168	1,042
(%)	(82.7)	(17.3)	(100.0)			
110～	189	40	229		32	261
(%)	(82.5)	(17.5)	(100.0)			
120～	75	20	95		11	106
(%)	(78.9)	(21.1)	(100.0)			
合計	23,624	4,592	28,216	72	5,456	33,744
(%)	(83.7)	(16.3)	(100.0)			
記載なし	3,778	898	4,676	12	1,514	6,202
(%)	(80.8)	(19.2)	(100.0)			
総計	27,402	5,490	32,892	84	6,970	39,946
(%)	(83.3)	(16.7)	(100.0)			
平均	75.46	75.08	75.40	75.75	75.59	75.43
標準偏差	12.92	13.44	13.01	12.42	12.55	12.93

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2014年調査